

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-272 796

**Use of Exposure Units for
Estimating Aquatic Toxicity
of Organophosphate Pesticides**

Environmental Research Lab, Duluth, Minn

Jul 77

SP4 085-3 17 517
Aug 1977

PB 272 796

Ecological Research Series



REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA 22161

TECHNICAL REPORT DATA (Please read instructions on the reverse of this form)		
1. REPORT NO. EPA-600/3-77-017	2. RECIPIENT'S OFFICE SYMBOL PA272796	
3. TITLE AND SUBTITLE USE OF EXPOSURE UNITS FOR ESTIMATING AQUATIC TOXICITY OF ORGANOPHOSPHATE PESTICIDES	4. REPORT DATE July 1977 Issuing date	5. PERFORMING ORGANIZATION CODE
6. AUTHOR(S) Donald T. Allison	7. PERFORMING ORGANIZATION REPORT NO.	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Environmental Research Laboratory - Duluth, MN Office of Research and Development U.S. Environmental Protection Agency Duluth, MN 55804	9. PROGRAM ELEMENT NO. 184-00	10. CONTRACT/GRANT NO.
11. SPONSORING AGENCY NAME AND ADDRESS SAME	12. TYPE OF REPORT AND PERIOD COVERED In-House	13. SPONSORING AGENCY CODE EPA/600/03
14. SUPPLEMENTARY NOTES		
15. ABSTRACT Environmental water concentrations of organophosphate pesticides can be expected to fluctuate widely due to use patterns and rapid hydrolysis. This study investigated some relationships of exposure concentration, duration and periodicity to the chronic toxicity of diazinon to flagfish (<i>Jordanella floridae</i>). Effects were compared on the basis of "exposure units" (exposure concentration x exposure duration). Treatments at equivalent exposure units caused similar overall effects on the test populations regardless of the life stages exposed or exposure duration per se. The hypothesis is proposed that exposure units could be used to estimate the environmental impact of fluctuating water concentrations of organophosphate pesticides over a wide range of concentration, duration, and periodicity.		
16. KEY WORDS AND DOCUMENT ANALYSIS		
17. DESCRIPTORS Pesticides* Organic phosphates* Diazinon* toxicity* Freshwater fishes	18. IDENTIFIERS/OPEN ENDED TERMS Exposure units Chronic toxicity Pulse exposure Flagfish	19. SUBST: Field/Group 6F 7C
20. DISTRIBUTION STATEMENT RELEASE TO PUBLIC	21. SECURITY CLASS (For Report) Unclassified 22. SECURITY CLASS (For Page) Unclassified	23. NO. OF PAGES 56 24. PRICE \$0.35 - \$0.1

EPA-600/3-77-077
July 1977

USE OF EXPOSURE UNITS FOR ESTIMATING
AQUATIC TOXICITY OF ORGANOPHOSPHATE PESTICIDES

by

Donald T. Allison
Environmental Research Laboratory-Duluth
Duluth, Minnesota 55804

ENVIRONMENTAL RESEARCH LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
DULUTH, MINNESOTA 55804

DISCLAIMER

This report has been reviewed by the Environmental Research Laboratory, Duluth, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

Our nation's fresh waters are vital for all animals and plants, yet our diverse uses of water--for recreation, food, energy, transportation, and industry--physically and chemically alter lakes, rivers, and streams. Such alterations threaten terrestrial organisms, as well as those living in water. The Environmental Research Laboratory in Duluth, Minnesota, develops methods, conducts laboratory and field studies, and extrapolates research findings

- to determine how physical and chemical pollution affects aquatic life
- to assess the effects of ecosystems on pollutants
- to predict effects of pollutants on large lakes through use of models
- to measure bioaccumulation of pollutants in aquatic organisms that are consumed by other animals, including man.

This report describes the results of a preliminary study of the relationships of exposure concentration, duration, and periodicity to the toxicity of an organophosphate pesticide. A hypothesis is proposed for estimation of aquatic environmental impact over a range of exposure conditions.

Donald I. Mount, Ph.D.
Director
Environmental Research Laboratory
Duluth, Minnesota

ABSTRACT

This study investigated the relationship of exposure duration, concentration, and periodicity with regard to the toxicity of one organophosphate insecticide to a freshwater fish. Flagfish (*Jordania floridae*) were exposed to uniform levels of diazinon throughout one and a half generations to establish baseline chronic toxicity values. Three additional tests were conducted with concentrations increased and duration decreased fourfold. These exposures were administered as a single pulse in each test to newly hatched, pre-spawning, and spawning flagfish, respectively. Subsequent effects on population success were observed until parents and progeny corresponded in age to the one and a half generations of the baseline test. To facilitate comparison, the toxicant challenges administered in each test were expressed as the product of exposure concentration and exposure duration ("exposure units").

All four types of exposure had similar overall effects on population growth and survival at equivalent exposure units. However, initiation of exposure during spawning caused temporary but complete inhibition of reproduction at concentrations which did not produce this effect in fish exposed continuously since hatch. This could have a severe impact on other species with a restricted reproductive period.

This study suggested the hypothesis that exposure units can be used in conjunction with previously established toxicity data to assess the environmental impact of fluctuating water concentrations of organophosphate pesticides over a wide range of concentration, duration, and periodicity.

CONTENTS

Foreword	144
Abstract	iv
Figures	vi
Tables	vii
Acknowledgment	viii
1. Introduction	1
2. Conclusions	3
3. Recommendations	4
4. Methods	5
Experimental design	5
Statistical analysis	5
5. Results	9
Terminal biomass in pulse and baseline chronic exposures	9
Survival of parental fish in pulse, chronic, and acute tests	9
6. Discussion	12
References	14
Appendix: Details of test techniques and results	15

FIGURES

<u>Number</u>		<u>Page</u>
1	Schematic profiles of the diazinon concentrations administered in the larval-juvenile, juvenile-adult, and adult-spawning pulse exposures	6
2	Schematic profiles of baseline-chronic and pulse exposure regimens	7
3	Estimated terminal biomass after one and a half generations expressed as percentage of control	10
4	Survival of parental fish to reproductive age expressed as percentage of control	11

TABLES

Number	Page
1 Chronic Effects on Flagfish (<i>Jordanella floridae</i>) of Continuous Life-Cycle Exposure to Dioxin	19
2 Relative Effects on Flagfish Progeny of Continuous Dioxin Exposure of Parents and Progeny from Paired Observations of Subdivided Egg Samples	20
3 Chronic Effects on Flagfish (<i>Jordanella floridae</i>) of Exposure During Larval-Juvenile Stage of Parents to 21-day Pulse of Dioxin	21
4 Chronic Effects on Flagfish (<i>Jordanella floridae</i>) of Exposure During Juvenile-Adult Stage of Parents to 21-day Pulse of Dioxin	22
5 Chronic Effects on Flagfish (<i>Jordanella floridae</i>) of Exposure During Adult-Spawning Stage of Parents to 21-day Pulse of Dioxin	23
6 Comparison of the Survival of Brook Trout and Fathead Minnows Versus Exposure Units of Dioxin	24

ACKNOWLEDGMENTS

The author wishes to thank Leonard H. Mueller and his assistants for diatom analyses; and John G. Eaton and other members of the Environmental Research Laboratory-Duluth for advice, assistance, and critical review of the manuscript.

SECTION I

INTRODUCTION

During the last decade many studies have been made of the long-term effects on aquatic organisms of continuous exposure to toxicants throughout complete or partial life cycles. However, pesticide concentrations in natural waters fluctuate widely because of use patterns (Bradshaw *et al.*, 1972) and variation in the physical, chemical, and biological characteristics of the water itself (Gomez *et al.*, 1969; Cowart *et al.*, 1971; Sethunathan and Pathak, 1972).

Periodicity of exposure seems to influence the toxicity of some pollutants. Acute toxicity of fluctuating levels of ammonia or zinc equaled that of constant-concentration exposure if the cycle times were short. Extending the cycle period increased the toxicity of ammonia but not of zinc (Brown *et al.*, 1969). Allison *et al.* (1963) reported accretion of tissue residues and progressive chronic toxicity over a period of 20 months in trout exposed for one-half hour every 4 weeks to near-lethal concentrations of DDT. A similar study by Allison (unpublished data) with the organophosphate pesticide malathion failed to produce conclusive chronic effects. However, continuous exposure to low levels of malathion will produce chronic effects (Munt and Stephan, 1967; Eaton, 1970).

The influence of duration and periodicity of exposure on chronic toxicity became a pertinent consideration following toxicity tests with diazinon (Allison and Hermanns, unpublished data). Continuous exposure caused chronic effects in fathead minnows and brook trout at concentrations less than 10^{-3} times their 96-hr LD_{50} 's. This great disparity makes the setting of a maximum permissible environmental level from the results of long-term continuous exposures to some toxicants unreasonable. A standard so derived would ignore the influence of exposure duration on the toxicity of a relatively non-persistent pesticide when the allowed maximum concentration probably would persist for only part of the life cycle of many aquatic species.

The transport, distribution, and persistence of organophosphate pesticides are subject to a large number of variable and generally unquantified factors. Estimates of exposure profiles under field conditions can be expected to have wide confidence limits. In addition, it would be impossible to simulate all probable field combinations of pesticide concentration and exposure duration in tests. Therefore, the current study was designed to compare in general the relative environmental impact of long-term, low-concentration exposure to a "non-persistent pesticide" with that of shorter term, higher concentration exposures intermediate between the currently standard acute and chronic toxicity tests. The possibility that major

differences might follow exposures limited to different life stages was included in the experimental design. It was also hoped that insight might be gained into any basic time-concentration relationship for the toxicity of organophosphate pesticides. However, no attempt was made to develop an experimental design capable of distinguishing and quantifying small differences for specific responses (growth, survival, egg hatch, etc. per ex).

The purposes of this paper are: (1) To note some problems inherent in assessments of the environmental hazards of "non-persistent" pesticides; (2) to present the method used to investigate the time-concentration relationship of organophosphate-pesticide toxicity; and (3) to present the hypothesis suggested by the results of the study.

SECTION 2

CONCLUSIONS

Similar overall response displayed at equivalent exposure units independent of the life stages exposed and exposure concentration or duration has led to the conclusion that use of exposure units might permit calculation of organophosphate pesticide toxicity over a wide range of water concentration, duration, and periodicity. Such a hypothesis would have important application to estimating the impact of organophosphate pesticides in practical field situations but it requires further testing and evaluation to verify its utility.

SECTION 3

RECOMMENDATIONS

The hypothesis that the exposure-unit concept can be used in estimating the impact of pesticides on aquatic populations rests on a limited data base. Therefore, the results of this study should be verified for other organophosphate pesticides, other species of fishes having longer egg-to-egg life spans, and aquatic invertebrates.

If acute toxicity tests are used to formulate water-quality criteria for short-term contamination by non-persistent pesticides, the possibility of residual chronic effects on survivors should be considered in addition to concurrent mortality.

SECTION 4

METHODS

EXPERIMENTAL DESIGN

Baseline chronic effects of several concentrations of the organophosphate insecticide diazinon on the flagfish (*Jordanella floridae*) were determined by continuous exposure throughout one and a half generations. The time factor involved was included by conversion to "exposure units" (exposure concentration $\times$ exposure duration) to permit comparison with the "challenge" experienced in exposures of shorter duration.

Three tests in which only a part of the life cycle was exposed used exposure units equivalent to the baseline chronic test (i.e., the same amount of diazinon used in each level of the baseline chronic were delivered in a shorter period of time by increasing concentration accordingly). These tests were designed so that the highest concentration experienced was about two-thirds the level known to be acutely toxic. To avoid shock from sudden exposure to near-acute concentrations and to approximate a profile of environmental contamination, maximum test concentrations were built up over a period of a few days and then decreased more slowly at termination of exposure. The result was a regimen in which concentrations were increased from zero to maximum in four equal daily increments. Maximum levels were maintained for 12 days and then reduced to zero in four equal decrements at 2-day intervals. Therefore, exposures in each test consisted of single pulses of diazinon delivered over a period of 21 days (Figure 1). These pulse exposures were administered to the separate test populations at 1, 29, and 66 days post hatch, respectively (Figure 2). The pulse tests were identified by the approximate life stage exposed as larval-juvenile, juvenile-adult, and adult-spawning. Post-exposure observations were made of the impact on treated parents and their untreated progeny to an age comparable to the baseline-chronic-test fish. In the adult-spawning test the progeny reared during exposure as well as the post-exposure progeny of treated parents were observed.

Details of test techniques are given in the appendix.

STATISTICAL ANALYSIS

The indices chosen to reflect population impact were parental survival and growth, egg production and hatch, and progeny survival and growth. The tests were run sequentially with different generations or spawnings from a common stock. Much of the extraneous variance inherent in sequential testing




Figure 1. Schematic profile of the diazinon concentrations administered in the larval-juvenile, juvenile-adult, and adult-spawning pulse exposures. ^aNominal concentration, ^bAverage measured concentrations in parentheses.




Figure 2. Schematic profiles of baseline-chronic and pulse exposure regimens. ^aOnly one set of equivalent exposure units is represented for each test.

was eliminated by expressing results as a percentage of the respective control. Comparison was further simplified by combining all data for specific effects into a common index of overall impact on population success which has been named terminal biomass. Details of the calculation of terminal biomass are given in the appendix.

SECTION 5

RESULTS

TERMINAL BIOMASS IN PULSE AND BASELINE-CHRONIC EXPOSURES

Detailed results used to calculate terminal biomass are presented in appendix, Tables 1, 3, 6, and 9.

Response regressions of terminal biomass for the baseline chronic and pulse tests are shown in Figure 3. As explained in the appendix, a time factor of 30 days was chosen to calculate exposure units in the baseline chronic test. For *Jordanella floridae* this time period included the life stages normally exposed in standard "life-cycle" tests with fishes. To show that a precise time value is not critical for demonstrating similar response between the pulse exposures and the life-cycle baseline exposure, the computational effects of a range of arbitrary baseline time frames between 60 and 120 days have also been delineated in Figure 3. The four types of exposure had similar impact on the test populations when treatment was quantified by exposure units.

SURVIVAL OF PARENTAL FISH IN PULSE, CHRONIC, AND ACUTE TESTS

In this study death of parental stock was limited to the actual exposure period in the pulse tests and to the pre-reproductive period in the baseline chronic test. This loss of breeding stock was a major factor influencing reproductive potential and estimates of terminal biomass. It seemed likely that deaths from acutely toxic exposure before the reproductive period would have a similar influence on population success. Calculations were therefore carried out to determine if exposure-unit relationship would hold true for exposures of short duration.

Four-day acute data were available for flagfish from a previous study (Allison and Hermanutz, unpublished data). Baseline chronic exposure units for comparison of parental deaths were based on 60 days of pre-reproductive exposure as no deaths occurred beyond this age. The regressions of parental survival for pulse, chronic, and acute exposures (Figure 4) indicate that correlation of effects with exposure units may have some validity in exposures as short as 4 days.

Calculations of parental survival in acute and chronic exposures of fathead minnow and brook trout were based on unpublished data from Allison and Hermanutz (appendix, Table 6). Although chronic data for survival below 75% were not available, there is some indication that the exposure-unit relationship may hold true for these species also.



Figure 3. Estimated terminal biomass after one and a half generations exposed to percentage of control.




Figure 4. Survival of juvenile fish to reproductive age expressed as percentage of control.

SECTION 6

DISCUSSION

This study was designed as a preliminary investigation of the possible relationship of exposure concentration, duration, and periodicity to the toxicity of organophosphate pesticides. If very different degrees of chronic response follow exposures of distinct life stages or intermittent versus continuous exposure, these factors should be considered when evaluating the environmental impact of non-persistent pesticides. The single-pulse exposure used here was intermediate to the acute and chronic exposures currently used to derive water-quality criteria. The pulse tests also investigated the impact on population success of exposure limited to three different periods in the life cycle of a freshwater fish.

Many of the specific effects accepted as significant in laboratory bioassays cannot be correlated with overall survival of a population. It was not the intent of this study to demonstrate differences in degree of sensitivity for specific observed effects. The objective was to determine if different patterns of exposure resulted in variations in impact on population success. This study did not indicate any extreme population sensitivity to any profile or timing of exposure. Instead response was correlated rather closely to the product of exposure concentration and exposure duration ("exposure units") over a wide range of time factors. However, there are obvious time and concentration constraints on the use of exposure units. Validity must at some point diminish on approaching very high or low concentrations for very short or long durations respectively.

Subjective evaluation would be required to determine the advisability of initiating exposure during a breeding season. Spawning was temporarily inhibited in the adult-spawning exposure (see appendix), but terminal biomass of Jordania was not greatly affected. However, if the physical and biological conditions necessary for successful reproduction in another species are of limited duration, an entire year class might be severely damaged by inhibition of spawning. Allison and Hermanutz (unpublished data) observed that brook trout, severely distressed for several weeks after initial exposure to diazinon, recovered and were capable of spawning after 6 months of continuous exposure. Although the continuous exposure did appear to cause some attrition, subjection of the trout to the same concentrations without acclimation just before their limited spawning period might have been more damaging to productivity.

In summary, the exposure-unit technique may have value in estimating the environmental impact of non-persistent pesticides over a range of exposure conditions. Estimated exposure units of a potential field situation could be summed and compared to exposure units of field or laboratory studies for which

results were already available. The concept should be verified with other species and toxicants and its limitations established and recognised before it is put to general use. In spite of restrictions it seems possible that the hypothesis may well be validated with sufficient confidence to warrant its inclusion in management decisions based on mass-balance modeling of water pollution.

SECTION 7

REFERENCES

- Allison, D., B. J. Saliman, and G. B. Gage. 1963. Insecticides, effects on cutthroat trout of repeated exposure to DDT. *Science* 142:958-961.
- Bradshaw, J. S., E. L. Loveridge, E. F. Rippes, J. L. Peterson, D. A. White, and D. E. Fuhrman. 1972. Seasonal variations in residues of chlorinated hydrocarbon pesticides in waters of the Utah Lake drainage system, 1970-1971. *West. Wildl. J.* 6:166-170.
- Brown, V. M., D. H. M. Jordan, and E. A. Tiller. 1969. The acute toxicity to rainbow trout of fluctuating concentrations and mixtures of ammonia, phenol and zinc. *J. Fish Biol.* 1:1-9.
- Cowart, R. P., F. L. Bonner, and E. A. Epps. 1971. Rate of hydrolysis of seven organophosphate pesticides. *Bull. Environ. Contam. Toxicol.* 6:231-236.
- Easton, J. G. 1970. Chronic malathion toxicity to bluegill (*Lepomis macrochirus* Rafinesque). *Water Res.* 4:673-684.
- Gomes, H. M., I. H. Suffet, and S. D. Faust. 1969. Kinetics of hydrolysis of diazinon and diaxozon. *Residue Rev.* 29:171-190.
- Mount, D. I., and G. E. Stephan. 1967. A method of establishing acceptable toxicant limits for fish -- malathion and the butoxysthanol ester of 2,4-D. *Trans. Am. Fish. Soc.* 96:185-193.
- Sethunathan, N., and M. D. Pathak. 1972. Increased biological hydrolysis of diazinon after repeated application in rice paddies. *J. Agr. Food Chem.* 20:585-589.

APPENDIX

DETAILS OF TEST TECHNIQUES AND RESULTS

METHODS

Baseline Chronic Exposure

Flagfish (*Jordanella floridae*) were used as the test species. Their relatively short life cycle (about 2 months egg-to-egg) allowed this series of tests to be conducted sequentially within a period of 2 years.

A modification of the proportional diluter equalized solvent concentrations in all test tanks. Temperatures were maintained between 25.5 and 26.5 C. Diazinon concentrations were nominally doubled between levels. Five concentrations (10-140 ug/l) and the control were duplicated. Exposure began with thirty 1-day-old flagfish per duplicate. Survival and growth were determined at 35 days and at termination. Each duplicate was thinned to two mature males and five mature females just prior to spawning. Two yarn-covered spawning substrates were placed in each duplicate chamber at 60 days, and spawning commenced immediately. After 60 days of spawning parental fish were killed and weighed.

Egg production was recorded daily. If numbers permitted, one sample of 30 eggs per duplicate was retained each day to measure hatch success. Twenty newly hatched larvae were placed in each of two larval chambers per duplicate on a space-available basis and were reared for 35 days to measure growth and survival (eight larval groups per treatment level).

Pulse Exposures

The pulse exposures were run sequentially in the test system used for the baseline chronic. The same basic test techniques were used. Fish initially exposed at 29 and 66 days (juvenile-adult and adult-spawning) were raised as pooled stock before testing. Observation of progeny growth and survival was limited to four groups per treatment level in the larval-juvenile and juvenile-adult tests as exposure had been discontinued before the reproductive period. Progeny produced early in the adult-spawning exposure (four groups per level) were maintained for 21 days before being terminated to make space available for post-exposure progeny studies of 35 days of growth and survival.

Chemical Analysis

Water levels of diazinon were monitored weekly in the baseline chronic and daily during the 21-day pulse exposures. Concentrations were determined by gas chromatography and electron capture detector.

Statistical Analysis

This study was conceived as a preliminary investigation to determine if exposures of different duration and seasonal occurrence might cause major differences in population success. Evidence of wide discrepancy might then warrant more sophisticated and costly experimental designs to quantify the factors involved. The possibility of extraneous variables (due to genetic and physical differences) as well as an inherent problem in the test design of precisely defining the time frame for each effect precluded direct comparison by standardized methods of statistical analysis and the use of mathematically derived tests of significance.

Exposure in the baseline chronic test lasted about 120 days. However, all deaths of parental fish took place before reproduction at 60 days. In addition, prolonged exposure of adults during reproduction apparently did not cause any correlated increase in effects on progeny. Therefore, for the purpose of comparison, 60 days was chosen as the exposure duration for parental deaths, and 90 days (60 days of parental exposure plus 30 days of progeny exposure) was chosen to represent a baseline chronic exposure for this species. The baseline chronic exposure period also encompassed the relative duration of a standard life-cycle exposure for fishes. The five levels of challenge could then be approximately quantified as "exposure units" to be duplicated in the pulse exposures. This same problem of defining the true exposure duration causing specific effects existed in the pulse exposures; consequently less confidence can be placed on the relationship of exposure units to specific effects than to overall impact on population from the entire exposure. Direct comparison and overall effects between these pulse tests is probably justified to the same extent as that usually made between different acute or chronic exposures with similar experimental designs. However, extrapolation of comparisons between baseline chronic, pulse, and acute exposures requires a greater degree of subjective evaluation.

The specific responses recorded in this study were parental survival and weight of survivors, egg production per spawning female and hatch success, and progeny survival and weight of survivors. These indices were used to derive an index of overall impact on population success, named terminal biomass. As a matter of convenience, terminal biomass for each treatment was computed and compared per each theoretical mating pair (one male and one female) present at the start of exposure. The average weights of parental males and females at test termination were multiplied by percentage parental survival, and the results were added to give an estimated index for parental biomass excluding the influence of thinning before reproduction. Progeny biomass was estimated by multiplying percentage parental survival to spawning by eggs per female by percentage hatch by percentage progeny survival to

termination by average weight of surviving progeny. The values for parental and progeny biomass were added for each treatment and expressed as a percentage of their respective test control. Extraneous variance due to sequential rather than simultaneous testing of the baseline and pulse exposures was thereby removed by equating all control results to 100%, and comparison could be made for population response versus exposure units.

Since actual exposure units delivered (determined by water analyses) differed somewhat between tests, comparison was made with calculated regressions of response. The regressions for the pulse tests are based on four data points, as the highest levels of exposure killed virtually all of the parental stock. Response within the range of the scaling plots presented in Figures 1 and 4 was fundamentally linear as represented.

Because true time factors in these tests could not be exactly defined, exposure-unit values must be considered as best-estimate approximations. Any tendency to assign unwarranted accuracy to the results should be avoided.

RESULTS

Specific results of the baseline chronic and 21-day pulse exposures are summarized in Tables 1 through 5 of this section. The response of progeny in the baseline test (Table 2) is of special interest. In previous tests with fathead minnows and brook trout (Allison and Hermanutz, unpublished data) all responses of the progeny at lower test concentrations could be attributed to the pre-spawning exposure of the parents. However, effects on flagfish progeny appear correlated to both parental and progeny exposure. In flagfish, progeny exposure alone apparently caused greater effects than parental exposure alone, but this evaluation is subjective because paired data could not be used in this case and analysis of unpaired data did not demonstrate a statistically significant difference. Diasticon also reduced the incubation time of flagfish eggs, an effect not seen with fathead minnows or brook trout.

Four samples of eggs per treatment were collected at the beginning of the adult-spawning test just before spawning was interrupted by the exposures. These eggs did not show any distinct evidence of reduced or early hatch as noted in the baseline chronic, even though the average concentrations experienced before hatch were generally higher than those to which eggs were continuously exposed in the baseline test. Graduated increases in concentration may have reduced toxicity, but more probably the sample size was simply too small to identify differences in response. During the remaining exposure period of 13 - 16 days following hatch of these eggs, the larvae displayed a range of response equivalent to that of fish of the same age exposed during the larval-juvenile pulse test.

In the highest concentration of each pulse test the fish were subjected to about two-thirds their 96-hr L₅₀ during the 12-day pulse peak. Sixty fish per test were exposed at this level, and only one of the 180 survived. Survival in lower exposures was adequate for comparison of growth, survival, and reproductive success. The three pulse tests did differ in degree of response for specific indices. For example, parental stock survived best in the larval-juvenile exposure and poorest in the juvenile-adult exposure.

Subsequent egg production per female was higher in the juvenile-adult exposure, but survival of progeny was poorer. These apparent differences may simply reflect variance due to different stocks of fish or slightly different physical conditions in sequential tests. In any event, the differences tended to cancel out during calculation of terminal biomass.

One effect was noted that was unique for the life stage exposed. Exposure in the adult-spawning pulse test was delayed until all groups commenced spawning. Spawning ceased abruptly when the various concentration profiles reached about 130 $\mu\text{g/l}$. Fish exposed to no more than 130 $\mu\text{g/l}$ for 12 days resumed spawning as soon as the concentration was reduced, but fish exposed to higher concentrations did not resume spawning until 1 - 2 weeks after concentrations were reduced below 130 $\mu\text{g/l}$. During the post-exposure period egg production in the treated groups equaled or exceeded the control. The temporary inhibition of spawning appears to be the result of initial exposure during the reproductive period. Fish in the baseline chronic exposed to about 240 $\mu\text{g/l}$ since hatch showed no evidence of spawning inhibition.

The actual impact of spawning inhibition on a fish population would depend on the length of the spawning season of a given species. In the flagfish the 21-day pulse exposure resulted in only a temporary interruption. Terminal biomass was calculated from equally weighted intra-exposure plus post-exposure reproductive data versus post-exposure data alone. Both methods of calculation resulted in biomass regressions within the range of those derived for exposure during the other two life stages.

CONCLUSIONS

THE UNIVERSITY OF CHICAGO

[illegible]

THE UNIVERSITY OF CHICAGO

TABLE 2. RELATIVE EFFECTS ON FLACFISH PROGENY OF CONTINUOUS DIAZINON EXPOSURE
OF PARENTS AND PROGENY FROM PAIRED OBSERVATIONS
OF SUBDIVIDED EGG SAMPLES

Concentration ($\mu\text{g/l}$)	Exposure of both parents and progeny VERSUS Exposure of parents only	Exposure of progeny only VERSUS No exposure of parents or progeny
<u>Start of hatch (days)</u>		
240	2.1 vs 2.7*	2.5 vs 3.5*
60	2.2 vs 2.2	2.5 vs 3.8*
<u>End of hatch (days)</u>		
240	3.6 vs 4.8*	4.0 vs 5.2*
60	3.3 vs 4.9*	4.4 vs 4.9*
<u>Hatch success</u>		
240	29% vs 46%	38% vs 62%
60	41% vs 52%	47% vs 59%
<u>Larval survival^a - hatch to 35 days</u>		
240	48% vs 92%	40% vs 75%
60	88% vs 92%	92% vs 78%
<u>Larval weight (μ)^a - 35 days post hatch</u>		
240	0.20 vs 0.26	0.19 vs 0.30
60	0.26 vs 0.27	0.24 vs 0.28

^aAverage of two subsamples per treatment.

*Pairs statistically different ($P \leq 0.05$) on the basis of "t" test. At least eight pairs were used for each comparison.

TABLE 1. CHRONIC EFFECTS ON PLACTIC (JEROME'S PLACTIC) OF EXPOSURE DURING

LARVAL--JUVENILE STAGES OF FISHES TO 11-200 MG/L OF DIOXIN^a

Exposure route (mg/l x days)	10,000	1,000	100	10	1	0	0
Group (1-day post-exposure (day))	10,000	1,000	100	10	1	0	0
<u>Survival Effects</u>							
Survived to 11 days post hatch	0	10	10	10	10	10	10
Average length at 11 days post hatch (mm)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Survived to day 100 (10 days post hatch) ^b	0	10	10	10	10	10	10
Average weight at 100 days post hatch (g)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Average weight of females at 100 days post hatch (g)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
<u>Reproductive Effects</u>							
24 production per female (mm)	1	1	1	1	1	1	1
Survival of eggs	1	1	1	1	1	1	1
<u>Adult Effects</u>							
Survived to 11 days post hatch	1	1	1	1	1	1	1
Average weight at 11 days post hatch (g)	1.0	1.0	1.0	1.0	1.0	1.0	1.0

^a Dioxin was added to water 1 and 11 days after hatch.^b Survival began 10 days after introduction of dioxin exposure.

Dioxin was added to water 1 and 11 days after hatch. 10,000 mg/l dioxin was added to water at hatch, 10,000 mg/l dioxin was added to water at 11 days after hatch.

TABLE 4. CRONIC EFFECTS OF FLUTER (*PHORAEA FLUTER*) OF EXPOSURE DURING JUVENILE-ADULT STAGE OF INDIAN TO 11-DEE PULSE OF FLUTER^a

Exposure effect (log ₁₀ + days)	11-DEE	11-DEE	11-DEE	11-DEE	11-DEE	11-DEE
average 11-DEE with concentration (log ₁₀)	1.10	1.10	1.10	1.10	1.10	1.10
Survival effect						
Survival to emergence 11 days post hatch ^b	0	0	0	0	0	0
average length 11 days post hatch (mm)	—	—	—	—	—	—
average weight 11 days post hatch (mg)	—	—	—	—	—	—
average weight at emergence 11 days post hatch (mg)	—	—	—	—	—	—
Mean weight effect						
log production per female parent	—	—	—	—	—	—
Survival to emergence 11 days post hatch	—	—	—	—	—	—
average weight 11 days post hatch (mg)	—	—	—	—	—	—
Survival effect						
Survival to emergence 11 days post hatch	—	—	—	—	—	—
average weight 11 days post hatch (mg)	—	—	—	—	—	—

^aSurvival effect exposed between 11 and 11 days after hatch.

^bSurvival to emergence 11 days after emergence of Indian exposure.

Significantly different from control (M.S.D.) using one-way analysis of variance and Student's *t*-test.

RECEIVED AT BOSTON MAY 12 10 30 AM 1862 BY EXPRESS OF THE NEW-YORK
 & ALBANY RAILROAD COMPANY

Year	Month	Day	Time	Place	Remarks
1900	Jan	1	10:00	San Francisco	Left for New York
1900	Jan	2	10:00	New York	Arrived
1900	Jan	3	10:00	New York	Left for New York
1900	Jan	4	10:00	New York	Arrived
1900	Jan	5	10:00	New York	Left for New York
1900	Jan	6	10:00	New York	Arrived
1900	Jan	7	10:00	New York	Left for New York
1900	Jan	8	10:00	New York	Arrived
1900	Jan	9	10:00	New York	Left for New York
1900	Jan	10	10:00	New York	Arrived
1900	Jan	11	10:00	New York	Left for New York
1900	Jan	12	10:00	New York	Arrived
1900	Jan	13	10:00	New York	Left for New York
1900	Jan	14	10:00	New York	Arrived
1900	Jan	15	10:00	New York	Left for New York
1900	Jan	16	10:00	New York	Arrived
1900	Jan	17	10:00	New York	Left for New York
1900	Jan	18	10:00	New York	Arrived
1900	Jan	19	10:00	New York	Left for New York
1900	Jan	20	10:00	New York	Arrived
1900	Jan	21	10:00	New York	Left for New York
1900	Jan	22	10:00	New York	Arrived
1900	Jan	23	10:00	New York	Left for New York
1900	Jan	24	10:00	New York	Arrived
1900	Jan	25	10:00	New York	Left for New York
1900	Jan	26	10:00	New York	Arrived
1900	Jan	27	10:00	New York	Left for New York
1900	Jan	28	10:00	New York	Arrived
1900	Jan	29	10:00	New York	Left for New York
1900	Jan	30	10:00	New York	Arrived
1900	Jan	31	10:00	New York	Left for New York

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

100

TABLE 6. SURVIVAL OF WHITE TROUT AND FATHEAD MINNOWS UNDER
EXPOSURE DATES OF 1960-1961^a

Fresh water			
1-10 exposure		11-20 exposure	
Exposure date	Survival	Exposure date	Survival
1-10	100	11-20	100
1-11	100	11-21	100
1-12	100	11-22	100
1-13	100	11-23	100
1-14	100	11-24	100
1-15	100	11-25	100
1-16	100	11-26	100
1-17	100	11-27	100
1-18	100	11-28	100
1-19	100	11-29	100
1-20	100	11-30	100
1-21	100	12-1	100
1-22	100	12-2	100
1-23	100	12-3	100
1-24	100	12-4	100
1-25	100	12-5	100
1-26	100	12-6	100
1-27	100	12-7	100
1-28	100	12-8	100
1-29	100	12-9	100
1-30	100	12-10	100
1-31	100	12-11	100
1-32	100	12-12	100
1-33	100	12-13	100
1-34	100	12-14	100
1-35	100	12-15	100
1-36	100	12-16	100
1-37	100	12-17	100
1-38	100	12-18	100
1-39	100	12-19	100
1-40	100	12-20	100
1-41	100	12-21	100
1-42	100	12-22	100
1-43	100	12-23	100
1-44	100	12-24	100
1-45	100	12-25	100
1-46	100	12-26	100
1-47	100	12-27	100
1-48	100	12-28	100
1-49	100	12-29	100
1-50	100	12-30	100
1-51	100	12-31	100
1-52	100	1-1	100
1-53	100	1-2	100
1-54	100	1-3	100
1-55	100	1-4	100
1-56	100	1-5	100
1-57	100	1-6	100
1-58	100	1-7	100
1-59	100	1-8	100
1-60	100	1-9	100
1-61	100	1-10	100
1-62	100	1-11	100
1-63	100	1-12	100
1-64	100	1-13	100
1-65	100	1-14	100
1-66	100	1-15	100
1-67	100	1-16	100
1-68	100	1-17	100
1-69	100	1-18	100
1-70	100	1-19	100
1-71	100	1-20	100
1-72	100	1-21	100
1-73	100	1-22	100
1-74	100	1-23	100
1-75	100	1-24	100
1-76	100	1-25	100
1-77	100	1-26	100
1-78	100	1-27	100
1-79	100	1-28	100
1-80	100	1-29	100
1-81	100	1-30	100
1-82	100	1-31	100
1-83	100	2-1	100
1-84	100	2-2	100
1-85	100	2-3	100
1-86	100	2-4	100
1-87	100	2-5	100
1-88	100	2-6	100
1-89	100	2-7	100
1-90	100	2-8	100
1-91	100	2-9	100
1-92	100	2-10	100
1-93	100	2-11	100
1-94	100	2-12	100
1-95	100	2-13	100
1-96	100	2-14	100
1-97	100	2-15	100
1-98	100	2-16	100
1-99	100	2-17	100
1-100	100	2-18	100
1-101	100	2-19	100
1-102	100	2-20	100
1-103	100	2-21	100
1-104	100	2-22	100
1-105	100	2-23	100
1-106	100	2-24	100
1-107	100	2-25	100
1-108	100	2-26	100
1-109	100	2-27	100
1-110	100	2-28	100
1-111	100	2-29	100
1-112	100	2-30	100
1-113	100	2-31	100
1-114	100	3-1	100
1-115	100	3-2	100
1-116	100	3-3	100
1-117	100	3-4	100
1-118	100	3-5	100
1-119	100	3-6	100
1-120	100	3-7	100
1-121	100	3-8	100
1-122	100	3-9	100
1-123	100	3-10	100
1-124	100	3-11	100
1-125	100	3-12	100
1-126	100	3-13	100
1-127	100	3-14	100
1-128	100	3-15	100
1-129	100	3-16	100
1-130	100	3-17	100
1-131	100	3-18	100
1-132	100	3-19	100
1-133	100	3-20	100
1-134	100	3-21	100
1-135	100	3-22	100
1-136	100	3-23	100
1-137	100	3-24	100
1-138	100	3-25	100
1-139	100	3-26	100
1-140	100	3-27	100
1-141	100	3-28	100
1-142	100	3-29	100
1-143	100	3-30	100
1-144	100	3-31	100
1-145	100	4-1	100
1-146	100	4-2	100
1-147	100	4-3	100
1-148	100	4-4	100
1-149	100	4-5	100
1-150	100	4-6	100
1-151	100	4-7	100
1-152	100	4-8	100
1-153	100	4-9	100
1-154	100	4-10	100
1-155	100	4-11	100
1-156	100	4-12	100
1-157	100	4-13	100
1-158	100	4-14	100
1-159	100	4-15	100
1-160	100	4-16	100
1-161	100	4-17	100
1-162	100	4-18	100
1-163	100	4-19	100
1-164	100	4-20	100
1-165	100	4-21	100
1-166	100	4-22	100
1-167	100	4-23	100
1-168	100	4-24	100
1-169	100	4-25	100
1-170	100	4-26	100
1-171	100	4-27	100
1-172	100	4-28	100
1-173	100	4-29	100
1-174	100	4-30	100
1-175	100	4-31	100
1-176	100	5-1	100
1-177	100	5-2	100
1-178	100	5-3	100
1-179	100	5-4	100
1-180	100	5-5	100
1-181	100	5-6	100
1-182	100	5-7	100
1-183	100	5-8	100
1-184	100	5-9	100
1-185	100	5-10	100
1-186	100	5-11	100
1-187	100	5-12	100
1-188	100	5-13	100
1-189	100	5-14	100
1-190	100	5-15	100
1-191	100	5-16	100
1-192	100	5-17	100
1-193	100	5-18	100
1-194	100	5-19	100
1-195	100	5-20	100
1-196	100	5-21	100
1-197	100	5-22	100
1-198	100	5-23	100
1-199	100	5-24	100
1-200	100	5-25	100
1-201	100	5-26	100
1-202	100	5-27	100
1-203	100	5-28	100
1-204	100	5-29	100
1-205	100	5-30	100
1-206	100	5-31	100
1-207	100	6-1	100
1-208	100	6-2	100
1-209	100	6-3	100
1-210	100	6-4	100
1-211	100	6-5	100
1-212	100	6-6	100
1-213	100	6-7	100
1-214	100	6-8	100
1-215	100	6-9	100
1-216	100	6-10	100
1-217	100	6-11	100
1-218	100	6-12	100
1-219	100	6-13	100
1-220	100	6-14	100
1-221	100	6-15	100
1-222	100	6-16	100
1-223	100	6-17	100
1-224	100	6-18	100
1-225	100	6-19	100
1-226	100	6-20	100
1-227	100	6-21	100
1-228	100	6-22	100
1-229	100	6-23	100
1-230	100	6-24	100
1-231	100	6-25	100
1-232	100	6-26	100
1-233	100	6-27	100
1-234	100	6-28	100
1-235	100	6-29	100
1-236	100	6-30	100
1-237	100	6-31	100
1-238	100	7-1	100
1-239	100	7-2	100
1-240	100	7-3	100
1-241	100	7-4	100
1-242	100	7-5	100
1-243	100	7-6	100
1-244	100	7-7	100
1-245	100	7-8	100
1-246	100	7-9	100
1-247	100	7-10	100
1-248	100	7-11	100
1-249	100	7-12	100
1-250	100	7-13	100
1-251	100	7-14	100
1-252	100	7-15	100
1-253	100	7-16	100
1-254	100	7-17	100
1-255	100	7-18	100
1-256	100	7-19	100
1-257	100	7-20	100
1-258	100	7-21	100
1-259	100	7-22	100
1-260	100	7-23	100
1-261	100	7-24	100
1-262	100	7-25	100
1-263	100	7-26	100
1-264	100	7-27	100
1-265	100	7-28	100
1-266	100	7-29	100
1-267	100	7-30	100
1-268	100	7-31	100
1-269	100	8-1	100
1-270	100	8-2	100
1-271	100	8-3	100
1-272	100	8-4	100
1-273	100	8-5	100
1-274	100	8-6	100
1-275	100	8-7	100
1-276	100	8-8	100
1-277	100	8-9	100
1-278	100	8-10	100
1-279	100	8-11	100
1-280	100	8-12	100
1-281	100	8-13	100
1-282	100	8-14	100
1-283	100	8-15	100
1-284	100	8-16	100
1-285	100	8-17	100
1-286	100	8-18	100
1-287	100	8-19	100
1-288	100	8-20	100
1-289	100	8-21	100
1-290	100	8-22	100
1-291	100	8-23	100
1-292	100	8-24	100
1-293	100	8-25	100
1-294	100	8-26	100
1-295	100	8-27	100
1-296	100	8-28	100
1-297	100	8-29	100
1-298	100	8-30	100
1-299	100	8-31	100
1-300	100	9-1	100
1-301	100	9-2	100
1-302	100	9-3	100
1-303	100	9-4	100
1-304	100	9-5	100
1-305	100	9-6	100
1-306	100	9-7	100
1-307	100	9-8	100
1-308	100	9-9	100
1-30			